Source - 1995

Source	*E. faecalis*	*E. faecium*	Other	Total	
Urine	893	36	21	950	(67%)
Other	283	17	7	307	(22%)
Blood/CSF	70	19	10	99	(7%)
Fluid	39	8	6	53	(4%)
All	1,285 (91%)	80 (6%)	44 (3%)	1,409	

Source - 1999

Source	*E. faecalis*	*E. faecium*	Other	Total	
Urine	1,080	102	9	1,191	(67%)
Other	326	37	6	369	(20%)
Blood/CSF	109	24	6	139	(8%)
Fluid	65	18	5	88	(5%)
All	1,580 (88%)	181 (10%)	26 (2%)	1,787	

As at November 2003

Source - 2003

Source	*E. faecalis*	*E. faecium*	Other	Total	
Urine	688	35	3	726	(67.2%)
Wound	89	4	2	95	(8.8%)
Blood/CSF	77	21	3	101	(9.3%)
Sterile cavity	55	10	1	66	(6.1%)
Other	83	7	1	91	(8.4%)
not stated	2			2	
All	994	77	10	1081	
	(92.0%)	(7.1%)	(0.9%)		

Enterococcus faecalis Resistance



As at November 2003

Australian Group on Antimicrobial Resistance

Enterococcus faecium Resistance



As at November 2003

Australian Group on Antimicrobial Resistance

High-level Aminoglycoside Resistance*

HLG	HLS	*E. faecalis* (n=642)		*E. faecium* (n=61)	
s	R	28	(4%)	9	(15%)
R	s	198	(31%)	8	(13%)
R	R	18	(3%)	21	(34%)
		244	(38%)	38	(62%)

* Where both gentamicin and streptomycin tested